TABL	E STEAT TO TELL	Page
C7.	Albemarle trimmed mean (ATM) data for all surface samples that are less than 2 standard deviations from the mean total population. The standard deviation, coefficient of variation, and the minimum and maximum concentration values used in this calculation for 22 elements (in $\mu g/g$ or ppm) in surface sediments of the Albemarle Sound estuarine system, are also included	
c8.		
D2.	estuarine system Summary of sediment particle size and the inorganic-organic composition of the total sediment for 356 samples of the	
D3.	Albemarle Sound estuarine system Summary of 350 samples and their inorganic and organic	
D4.	composition within the Albemarle Sound estuarine system Changing patterns of % sand, mud, and organic matter in	
D5.	samples through the Albemarle Sound estuarine system	
D6.	estuarine system	
D7.	Average composition of total sediment for all samples collected within the entire Pamlico, Neuse, and Albemarle	
D8.	estuarine systems and within the respective trunk estuaries Comparison of trimmed mean concentrations for 22 elements (in $\mu g/g$ or ppm) in surface sediments of Albemarle Sound (this study), the Neuse River (Riggs et al., 1991c), and	93
	Pamlico River (Riggs et al., 1989b) estuarine systems	94
E1.	Outline of the regional areas of the Albemarle Sound estuarine system and associated figures and tables	
E2.	considered in the Results	
E3.	samples collected in Welch Creek	
E4.	samples collected in the Lower Roanoke River Summary of mean and maximum enrichment factors for 15 trace elements in surface sediments from Middle River and Cashie	
	River, two tributary channels of the Lower Roanoke River	106